

References

- [1] Bang, H., & Dyerberg, J. (1972). Plasma lipids and lipoproteins in Greenlandic west coast Eskimos. *Acta Medica Scandinavica*, 192(1-6), 85-94.
- [2] Bang, H., Dyerberg, J., & Nielsen, A. n. (1971). Plasma lipid and lipoprotein pattern in Greenlandic West-coast Eskimos. *The Lancet*, 297(7710), 1143-1146.
- [3] Kris-Etherton, P. M., Harris, W. S., & Appel, L. J. (2002). Fish consumption, fish oil, omega-3 fatty acids, and cardiovascular disease. *Circulation*, 106(21), 2747-2757.
- [4] Adkins, Y., & Kelley, D. S. (2010). Mechanisms underlying the cardioprotective effects of omega-3 polyunsaturated fatty acids. *The Journal of nutritional biochemistry*, 21(9), 781-792.
- [5] Sekikawa, A., Cui, C., Sugiyama, D., Fabio, A., Harris, W. S., & Zhang, X. (2019). Effect of High-Dose Marine Omega-3 Fatty Acids on Atherosclerosis: A Systematic Review and Meta-Analysis of Randomized Clinical Trials. *Nutrients*, 11(11), 2599.
- [6] Davidson, M. H. (2006). Mechanisms for the hypotriglyceridemic effect of marine omega-3 fatty acids. *The American journal of cardiology*, 98(4), 27-33.
- [7] Wei, M. Y., & Jacobson, T. A. (2011). Effects of eicosapentaenoic acid versus docosahexaenoic acid on serum lipids: a systematic review and meta-analysis. *Current atherosclerosis reports*, 13(6), 474-483.
- [8] Lopez-Huertas, E. (2012). The effect of EPA and DHA on metabolic syndrome patients: a systematic review of randomised controlled trials. *British journal of nutrition*, 107(S2), S185-S194.
- [9] Allaire, J., Couture, P., Leclerc, M., Charest, A., Marin, J., Lépine, M.-C., . . . Lamarche, B. (2016). A randomized, crossover, head-to-head comparison of eicosapentaenoic acid and docosahexaenoic acid supplementation to reduce inflammation markers in men and women: the Comparing EPA to DHA (ComparED) Study. *The American journal of clinical nutrition*, 104(2), 280-287.
- [10] Calder, P. C. (2006). n- 3 polyunsaturated fatty acids, inflammation, and inflammatory diseases. *The American journal of clinical nutrition*, 83(6), 1505S-1519S.
- [11] Calder, P. C. (2011). Fatty acids and inflammation: the cutting edge between food and pharma. *European journal of pharmacology*, 668, S50-S58.
- [12] Calder, P. C., & Yaqoob, P. (2009). Understanding omega-3 polyunsaturated fatty acids. *Postgraduate medicine*, 121(6), 148-157.

- [13] Calder, P. C. (2014). Very long chain omega-3 (n-3) fatty acids and human health. *European journal of lipid science and technology*, 116(10), 1280-1300.
- [14] Ghasemifard, S., Turchini, G. M., & Sinclair, A. J. (2014). Omega-3 long chain fatty acid "bioavailability": A review of evidence and methodological considerations. *Progress in lipid research*, 56, 92-108.
- [15] Zárate, R., el Jaber-Vazdekis, N., Tejera, N., Pérez, J. A., & Rodríguez, C. (2017). Significance of long chain polyunsaturated fatty acids in human health. *Clinical and translational medicine*, 6(1), 25.
- [16] Thesing, C. S., Bot, M., Milaneschi, Y., Giltay, E. J., & Penninx, B. W. (2018). Omega-3 polyunsaturated fatty acid levels and dysregulations in biological stress systems. *Psychoneuroendocrinology*, 97, 206-215.
- [17] Gorjão, R., Azevedo-Martins, A. K., Rodrigues, H. G., Abdulkader, F., Arcisio-Miranda, M., Procopio, J., & Curi, R. (2009). Comparative effects of DHA and EPA on cell function. *Pharmacology & therapeutics*, 122(1), 56-64.
- [18] Stillwell, W., & Wassall, S. R. (2003). Docosahexaenoic acid: membrane properties of a unique fatty acid. *Chemistry and physics of lipids*, 126(1), 1-27.
- [19] Hashimoto, M., Hossain, S., Al Mamun, A., Matsuzaki, K., & Arai, H. (2017). Docosahexaenoic acid: one molecule diverse functions. *Critical reviews in biotechnology*, 37(5), 579-597.
- [20] Christensen, J. H. (2011). Omega-3 polyunsaturated fatty acids and heart rate variability. *Frontiers in physiology*, 2, 84.
- [21] Christensen, J. H., & Schmidt, E. B. (2007). Autonomic nervous system, heart rate variability and n-3 fatty acids. *Journal of Cardiovascular Medicine*, 8, S19-S22.
- [22] Katan, M. B., Deslypere, J., Van Birgelen, A., Penders, M., & Zegwaard, M. (1997). Kinetics of the incorporation of dietary fatty acids into serum cholesteryl esters, erythrocyte membranes, and adipose tissue: an 18-month controlled study. *Journal of lipid research*, 38(10), 2012-2022.
- [23] Cao, J., Schwichtenberg, K. A., Hanson, N. Q., & Tsai, M. Y. (2006). Incorporation and clearance of omega-3 fatty acids in erythrocyte membranes and plasma phospholipids. *Clinical chemistry*, 52(12), 2265-2272.
- [24] Browning, L. M., Walker, C. G., Mander, A. P., West, A. L., Madden, J., Gambell, J. M., . . . Calder, P. C. (2012). Incorporation of eicosapentaenoic and docosahexaenoic acids into lipid pools when given as supplements providing doses equivalent to typical intakes of oily fish. *The American journal of clinical nutrition*, 96(4), 748-758.

- [25] Harris, W. S., & Thomas, R. M. (2010). Biological variability of blood omega-3 biomarkers. *Clinical biochemistry*, 43(3), 338-340.
- [26] Brenna, J. T., Plourde, M., Stark, K. D., Jones, P. J., & Lin, Y.-H. (2018). Best practices for the design, laboratory analysis, and reporting of trials involving fatty acids. *The American journal of clinical nutrition*, 108(2), 211-227.
- [27] Ghasemi Fard, S., Wang, F., Sinclair, A. J., Elliott, G., & Turchini, G. M. (2018). How does high DHA fish oil affect health? A systematic review of evidence. *Critical reviews in food science and nutrition*, 59(11), 1684-1727.
- [28] Meyer, B. J., & De Groot, R. H. (2017). Effects of omega-3 long chain polyunsaturated fatty acid supplementation on cardiovascular mortality: the importance of the dose of DHA. *Nutrients*, 9(12), 1305.
- [29] Lawson, L. D., & Hughes, B. G. (1988). Human absorption of fish oil fatty acids as triacylglycerols, free acids, or ethyl esters. *Biochemical and biophysical research communications*, 152(1), 328-335.
- [30] Zuijdgeest-van Leeuwen, S. D., Dagnelie, P. C., Rietveld, T., van den Berg, J. W. O., & Wilson, J. P. (1999). Incorporation and washout of orally administered n-3 fatty acid ethyl esters in different plasma lipid fractions. *British journal of nutrition*, 82(6), 481-488.
- [31] Neubronner, J., Schuchardt, J. P., Kressel, G., Merkel, M., von, von Schacky, C., & Hahn, A. (2011). Enhanced increase of omega-3 index in response to long-term n-3 fatty acid supplementation from triacylglycerides versus ethyl esters. *European journal of clinical nutrition*, 65(2), 247-254.
- [32] Trabulsi, J., & Schoeller, D. A. (2001). Evaluation of dietary assessment instruments against doubly labeled water, a biomarker of habitual energy intake. *American journal of physiology-endocrinology and metabolism*, 281(5), E891-E899.
- [33] Engel, M. G., Kern, H. J., Brenna, J. T., & Mitmesser, S. H. (2018). Micronutrient gaps in three commercial weight-loss diet plans. *Nutrients*, 10(1), 108.
- [34] Bailey-Hall, E., Nelson, E. B., & Ryan, A. S. (2008). Validation of a rapid measure of blood PUFA levels in humans. *Lipids*, 43(2), 181-186.
- [35] Bell, J. G., Mackinlay, E. E., Dick, J. R., Younger, I., Lands, B., & Gilhooly, T. (2011). Using a fingertip whole blood sample for rapid fatty acid measurement: method validation and correlation with erythrocyte polar lipid compositions in UK subjects. *British journal of nutrition*, 106(9), 1408-1415.
- [36] Gunash, J., Aristizabal-Henao, J. J., & Stark, K. D. (2019). Quantitating fatty acids in dried blood spots on a common collection card versus a novel wicking sampling device. *Prostaglandins, leukotrienes and essential fatty acids*, 145, 1-6.

- [37] Harris, W., & Polreis, J. (2016). Measurement of the omega-3 index in dried blood spots. *Ann. Clin. Lab. Res*, 4(4).
- [38] Lin, Y. H., Hanson, J. A., Strandjord, S. E., Salem, N. M., Dretsch, M. N., Haub, M. D., & Hibbeln, J. R. (2014). Fast transmethylation of total lipids in dried blood by microwave irradiation and its application to a population study. *Lipids*, 49(8), 839-851.
- [39] Liu, G., Mühlhäusler, B. S., & Gibson, R. A. (2014). A method for long term stabilisation of long chain polyunsaturated fatty acids in dried blood spots and its clinical application. *Prostaglandins, leukotrienes and essential fatty acids*, 91(6), 251-260.
- [40] Marangoni, F., Colombo, C., & Galli, C. (2004). A method for the direct evaluation of the fatty acid status in a drop of blood from a fingertip in humans: applicability to nutritional and epidemiological studies. *Analytical biochemistry*, 326(2), 267-272.
- [41] Marangoni, F., Colombo, C., Martiello, A., Negri, E., & Galli, C. (2007). The fatty acid profiles in a drop of blood from a fingertip correlate with physiological, dietary and lifestyle parameters in volunteers. *Prostaglandins, leukotrienes and essential fatty acids*, 76(2), 87-92.
- [42] Metherel, A., Armstrong, J., Patterson, A., & Stark, K. (2009). Assessment of blood measures of n-3 polyunsaturated fatty acids with acute fish oil supplementation and washout in men and women. *Prostaglandins, leukotrienes and essential fatty acids*, 81(1), 23-29.
- [43] Metherel, A. H., Buzikievich, L. M., Charkhzarin, P., Patterson, A. C., Peel, A. C., Howorth, A. M., . . . Stark, K. D. (2012). Omega-3 polyunsaturated fatty acid profiling using fingertip-prick whole blood does not require overnight fasting before blood collection. *Nutrition Research*, 32(8), 547-556.
- [44] Metherel, A. H., Hogg, R. C., Buzikievich, L. M., & Stark, K. D. (2013). Butylated hydroxytoluene can protect polyunsaturated fatty acids in dried blood spots from degradation for up to 8 weeks at room temperature. *Lipids in health and Disease*, 12(1), 22.
- [45] Metherel, A. H., & Stark, K. D. (2016). The stability of blood fatty acids during storage and potential mechanisms of degradation: a review. *Prostaglandins, leukotrienes and essential fatty acids*, 104, 33-43.
- [46] Min, Y., Ghebremeskel, K., Geppert, J., & Khalil, F. (2011). Effect of storage temperature and length on fatty acid composition of fingertip blood collected on filter paper. *Prostaglandins, leukotrienes and essential fatty acids*, 84(1-2), 13-18.
- [47] Simopoulos, A. P. (2006). Evolutionary aspects of diet, the omega-6/omega-3 ratio and genetic variation: nutritional implications for chronic diseases. *Biomedicine & pharmacotherapy*, 60(9), 502-507.

- [48] Blasbalg, T. L., Hibbeln, J. R., Ramsden, C. E., Majchrzak, S. F., & Rawlings, R. R. (2011). Changes in consumption of omega-3 and omega-6 fatty acids in the United States during the 20th century. *The American journal of clinical nutrition*, 93(5), 950-962.
- [49] Simopoulos, A. P. (2004). Omega-6/omega-3 essential fatty acid ratio and chronic diseases. *Food reviews international*, 20(1), 77-90.
- [50] Kuratko, C. N., & Salem Jr, N. (2009). Biomarkers of DHA status. *Prostaglandins, leukotrienes and essential fatty acids*, 81(2-3), 111-118.